

Embargoed until Monday, 7 April 2008, 17:00 GMT

An unexpected way to cause leukaemia

New mouse model grants insight into the genetic and molecular mechanisms underpinning acute myeloid leukaemia

Monterotondo, 8 April 2008 - Leukaemia – cancer of blood or bone marrow – is caused by mutations that allow defective blood cells to accumulate and displace healthy blood. To devise effective therapies it is crucial to know which mutations cause leukaemia and which cell type gives rise to leukaemic cells. Researchers from the European Molecular Biology Laboratory (EMBL) in Italy, the EMBL-European Bioinformatics Institute, UK, and the Universities of Harvard, USA, and Lund, Sweden, have now used genetic engineering to introduce a mutation found in human leukaemia patients into mice. In the current issue of *Cancer Cell* they report that the mutation causes leukaemia by triggering innate genetic programmes that allow white blood cells to proliferate uncontrollably. The findings have implications for the way leukaemia should be treated.

Blood is generated from a small number of multipotent stem cells that divide, differentiate and give rise to the many different cell types that make up the blood. At the same time they also maintain the pool of stem cells through a process called self-renewal. While differentiating, cells acquire specific properties and functions, but lose the capacity to self-renew in the way stem cells do. Mutations interfering with this process and promoting uncontrolled proliferation of certain blood cells can lead to leukaemia. Researchers of the group of Claus Nerlov at EMBL's Mouse Biology Unit now prove that a mutation in a protein called C/EBP α causes acute myeloid leukaemia (AML), a type of leukaemia affecting one lineage of white blood cells, in mice.

“10 percent of all patients suffering from AML have this mutation, but we could never be sure if it causes the disease. By precisely reproducing the human mutation in the mouse we now proved a causative relation,” says Peggy Kirstetter, who carried out the research in Nerlov's lab.

Instead of promoting uncontrolled proliferation of malignant blood stem cells, as often assumed as the cause of leukaemia, the mutation acts on already partially differentiated cells. It reprogrammes these cells to self-renew and to produce countless dysfunctional daughter cells, which displace the healthy blood cells, eventually leading to the inability to transport oxygen around in the body.

“This is the first time that non-stem cell myeloid leukaemia has been generated within a healthy blood system. The findings will have profound implications for our understanding of the development and treatment of leukaemias,” says Nerlov.

Scientists always thought that the mutation was the crucial step leading to leukaemia that should be targeted by drugs. Nerlov and colleagues identified a genetic program activated in self-renewing leukemic cells, which is shared with similar leukaemias caused by other types of mutations. The findings suggest that the cellular changes that lead to self-renewal are mutation-independent. To develop drugs with a more general efficacy it may therefore be more efficient to target the molecules and pathways shared between different cancer stem cells.

Source Article

P. Kirstetter, M.B. Schuster, O. Bereshchenko, S. Moore, H. Dvinge, E. Kurz, K. Theilgaard-Mönch, R. Månsson, T.Å. Pedersen, T. Pabst, E. Schrock, B.T. Porse, S.E.W. Jacobsen, P. Bertone, D.G. Tenen & C. Nerlov. Modeling of C/EBP α mutant acute myeloid leukemia reveals a common expression signature of committed myeloid leukemia initiating cells. *Cancer Cell*, 8 April 2008

Contact:

Anna-Lynn Wegener, EMBL Press Officer, Heidelberg, Germany, Tel: +49 6221 387 452, www.embl.org, wegener@embl.de

About EMBL

The European Molecular Biology Laboratory is a basic research institute funded by public research monies from 20 member states (Austria, Belgium, Croatia, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Israel, Italy, Luxembourg, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and the United Kingdom) and associate member state Australia. Research at EMBL is conducted by approximately 80 independent groups covering the spectrum of molecular biology. The Laboratory has five units: the main Laboratory in Heidelberg, and Outstations in Hinxton (the European Bioinformatics Institute), Grenoble, Hamburg, and Monterotondo near Rome. The cornerstones of EMBL's mission are: to perform basic research in molecular biology; to train scientists, students and visitors at all levels; to offer vital services to scientists in the member states; to develop new instruments and methods in the life sciences and to actively engage in technology transfer activities. EMBL's International PhD Programme has a student body of about 170. The Laboratory also sponsors an active Science and Society programme. Visitors from the press and public are welcome.

Policy regarding use

EMBL press releases may be freely reprinted and distributed via print and electronic media. Text, photographs & graphics are copyrighted by EMBL. They may be freely reprinted and distributed in conjunction with this news story, provided that proper attribution to authors, photographers and designers is made. High-resolution copies of the images can be downloaded from the EMBL web site: www.embl.org